Adult Patient Vaccine Tracker

Saving Lives: Integrating Vaccines for Adults Into Routine Care

Use with the Recommended Adult Immunization Schedule from the Centers for Disease Control and Prevention on the reverse side of this document.

Patient Name:	
Date of Birth:	
Chart Number:	

Vaccine:	Vaccine Name/Type	Site	Date Given: (mm/dd/yyyy)	Vaccine		Date on VIS	HCP Initials
				Lot #	MFR.		
Influenza							
HPV			1.				
			2.				
			3.				
Zoster							
Tdap							
Td			1.				
			2.				
			3.				
Pneumococcal			1.				
			2.				
			3.				
Meningococcal			1.				
			2.				
Varicella							
MMR			1.				
			2.				
Hepatitis A			1.				
			2.				
Hepatitis B			1.				
			2.				
			3.				

National Foundation for Infectious Diseases®

This document is supported by an unrestricted educational grant from Pfizer Inc. The National Foundation for Infectious Diseases' (NFID) policies prohibit funders from controlling program content. To view NFID's policy for unrestricted educational grants, visit www.nfid.org/info/grantspolicy.pdf.

December 2012

2012 Adult Immunization Schedule

For use with the Adult Patient Immunization Tracker on the reverse side of this document.
Source: Centers for Disease Control and Prevention (CDC)

Recommended Adult Immunization Schedule

UNITED STATES · 2012

Note: These recommendations *must* be read with the footnotes that follow containing number of doses, intervals between doses, and other important information.

Recommended adult immunization schedule, by vaccine and age group

VACCINE ▼ AGE GROUP ►	19-21 years	22-26 years	27-49 years	50-59 years	60-64 years	≥ 65 years
Influenza [2]	1 dose annually					
Tetanus, diphtheria, pertussis (Td/Tdap) [3,*]	Substitute 1-time dose of Tdap for Td booster; then boost with Td every 10 yrs					Td/Tdap[3]
Varicella [4,*]	2 Doses					
Human papillomavirus (HPV) Female [5,*]	3 doses					
Human papillomavirus (HPV) Male [5,*]	3 doses					
Zoster [6]					1 dose	
Measles, mumps, rubella (MMR) [7,*]	1 or 2 doses			1 dose		
Pneumococcal (polysaccharide) [8,9]	1 or 2 doses					1 dose
Meningococcal [10,*]	1 or more doses					
Hepatitis A [11,*]	2 doses					
Hepatitis B [12,*]	3 doses					

*Covered by the Vaccine Injury Compensation Program

- For all persons in this category who meet the age requirements and who lack documentation of vaccination or have no evidence of previous infection
- Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)
- Tdap recommended for ≥65 if contact with <12 month old child. Either Td or Tdap can be used if no infant contact
- No recommendation

Report all clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System (VAERS). Reporting forms and instructions on filing a VAERS report are available at www.vaers.hhs.gov or by telephone, 800-822-7967.

Information on how to file a Vaccine Injury Compensation Program claim is available at www.hrsa.gov/vaccinecompensation or by telephone, 800-338-2382. To file a claim for vaccine injury, contact the U.S. Court of Federal Claims, 717 Madison Place, N.W., Washington, D.C. 20005; telephone, 202-357-6400.

Additional information about the vaccines in this schedule, extent of available data, and contraindications for vaccination is also available at www.cdc.gov/vaccines or from the CDC-INFO Contact Center at 800-CDC-INFO (800-232-4636) in English and Spanish, 8:00 a.m. - 8:00 p.m. Eastern Time, Monday - Friday, excluding holidays.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

Vaccines that might be indicated for adults based on medical and other indications

VACCINE ▼ INDICATION ►	Pregnancy	Immunocompromising conditions (excluding human immunodeficiency virus [HIV])[4,6,7,14]	HIV infection[4,7,13,14] CD4+ T lymphocyte count <200 cells/μL	HIV infection[4,7,13,14] CD4+ T lymphocyte count >200 cells/μL	Men who have sex with men (MSM)	Heart disease, chronic lung disease, chronic alcoholism	Asplenia[13] (including elective splenectomy and persistent complement component deficiencies)	Chronic liver disease	Diabetes, kidney failure, end-stage renal disease, receipt of hemodialysis	Health-care personnel
Influenza [2]	1 dose TIV annually				1 dose TIV or LAIV annually	1 dose TIV annually				1 dose TIV or LAIV annually
Tetanus, diphtheria, pertussis (Td/Tdap) [3,*]	Substitute 1-time dose of Tdap for Td booster; then boost with Td every 10 yrs									
Varicella [4,*]	Contraindicated			2 doses						
Human papillomavirus (HPV) Female [5,*]		3 doses through age 26 yrs				3 doses through age 26 yrs				
Human papillomavirus (HPV) Male [5,*]		3 doses through age 26 yrs				3 doses through age 21 yrs				
Zoster [6]	Contraindicated				1 dose					
Measles, mumps, rubella (MMR) [7,*]	Contraindicated			1 or 2 doses						
Pneumococcal (polysaccharide) [8,9]	1 or 2 doses									
Meningococcal [10,*]	1 or more doses									
Hepatitis A [11,*]	2 doses									
Hepatitis B [12,*]	3 doses									

*Covered by the Vaccine Injury Compensation Program

- For all persons in this category who meet the age requirements and who lack documentation of vaccination or have no evidence of previous infection
- Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)
- Contraindicated
- No recommendation

The recommendations in this schedule were approved by the Centers for Disease Control and Prevention's (CDC) Advisory Committee on Immunization Practices (ACIP), the American Academy of Family Physicians (AAFP), the American College of Physicians (ACP), American College of Obstetricians and Gynecologists (ACOG) and American College of Nurse-Midwives (ACNM).

These schedules indicate the recommended age groups and medical indications for which administration of currently licensed vaccines is commonly indicated for adults ages 19 years and older, as of January 1, 2012. For all vaccines being recommended on the Adult Immunization Schedule: a vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Licensed combination vaccines may be used whenever any components of the combination are indicated and when the vaccine's other components are not contraindicated. For detailed recommendations on all vaccines, including those used primarily for travelers or that are issued during the year, consult the manufacturers' package inserts and the complete statements from the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/pubs/acip-list.htm). Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

Since the 2012 Adult Immunization Schedule was published in February 2012, CDC has made updated adult recommendations for pneumococcal disease. For more information, please visit: http://adultvaccination.org/professional-resources/pneumo-toolkit/adult-pneumo-guide-hcp.pdf.

NFID Tips:

- CDC publishes the Recommended Adult Immunization Schedule annually with footnotes for additional guidance. Visit cdc.gov/vaccines for the most current schedule and accompanying footnotes.
- Print the tracker on cardstock and place a self-sticking file tab on the edge of the page for easy identification. Attach tracker to inside front or back of patient's file.
- Print this schedule on the reverse side (or staple as page two) of the Adult Patient Vaccine Tracker for easy reference.
- Screen patients for vaccination needs upon check-in.
- Talk to patients about vaccines at all office visits (well, sick, follow-up).
- Ask patients if they have received any vaccinations at other places (eg, another medical office, pharmacy, public health clinic, place of employment, etc.).
- Encourage patients to learn about vaccination. Educational resources and template practice communications are available on **Adultvaccination.org.**